

# New Results on the Theoretical Precision of the LEP/SLC Luminosity\*

S. Jadach<sup>†</sup>, M. Melles<sup>‡</sup>, B.F.L. Ward, and S. A. Yost  
*Department of Physics and Astronomy, University of Tennessee,  
Knoxville, Tennessee 37996-1200, USA*

May 1998  
UTHEP-98-0501

ICHEP98, VANCOUVER, JULY, 1998

## OUTLINE

- INTRODUCTION
- EXACT RESULTS ON  $O(\alpha)$  CORRECTION TO  $1\gamma$  BREMSSTRAHLUNG
- EXACT RESULTS ON  $2\gamma$  BREMSSTRAHLUNG: TECHNICAL PRECISION CHECK
- EXACT 2-LOOP VIRTUAL CORRECTION
- NEW ERROR ANALYSIS FOR BHLUMI 4.04
- CONCLUSIONS

# ● INTRODUCTION

## \* MOTIVATION

### PRECISION SM TESTS AT LEP

$$\Delta\sigma/\sigma = \left( \left( \frac{\partial\sigma}{\partial\alpha} \left( \frac{\Delta\alpha}{\alpha} \right) \frac{\sigma}{\sigma} \right)^2 + \dots \right)^{1/2}$$

⇒  $\frac{\Delta\alpha}{\alpha}$  NEEDED AS SMALL AS POSSIBLE

⇒  $\frac{\Delta\alpha}{\alpha} \Big|_{th}$  NEEDED AS SMALL AS POSSIBLE

## \* STATUS

$\frac{\Delta\sigma}{\sigma} \Big|_{TH}$

(A.P.P. B28, 925  
(1997))

Type of correction/error	LEP1		LEP2
	Past	Present	Present
(a) Missing photonic $O(\alpha^2 L)$	0.15%	0.10%	0.20%
(b) Missing photonic $O(\alpha^3 L^3)$	0.008%	0.015%	0.03%
(c) Vacuum polarization	0.05%	0.04%	0.10%
(d) Light pairs	0.01%	0.03%	0.05%
(e) Z-exchange	0.03%	0.015%	0.0%
Total	0.16%	0.11%	0.25%

Table 1: Summary of the total (physical+technical) theoretical uncertainty for a typical calorimetric detector. For LEP1, the above estimate is valid for the angular range  $1^\circ$ - $3^\circ$ ; for LEP2 it covers energies up to 176 GeV, and angular ranges of  $1^\circ$ - $3^\circ$  and  $3^\circ$ - $6^\circ$  (see the text for further comments).

$\frac{\Delta\sigma}{\sigma} \Big|_{EXPT} \lesssim .05\%$

⇒ NEED TO MOVE  $\frac{\Delta\sigma}{\sigma}$  TO

.05%-REGIME, AT LEAST.

\* DOMINANT CONTRIBUTION TO  $\Delta\sigma/\sigma|_{TH}$

(a) MISSING PHOTONIC  $O(\alpha^2 L) \leftrightarrow .1\%$

● PHYSICAL PRECISION

1.  $O(\alpha)$  CORR. TO  $1\gamma$  BREMSSTRAHLUNG:  
 $\bar{\beta}_1$

2.  $O(\alpha^2)$  CORR. TO  $\bar{\beta}_0$  (2-LOOP  
VIRTUAL CORR.)

● TECHNICAL PRECISION


1.  $O(\alpha^2)$   $2\gamma$  BREMSSTRAHLUNG:  
 $\bar{\beta}_2$

(INDEPENDENT REALIZATION OF  
 $2\gamma$  PHASE SPACE IN 'TEST'  
MC)

WE NOW TURN TO THESE IN TURN

(4)  
● EXACT RESULTS ON  $O(\alpha)$  CORRECTION  
TO  $1\gamma$  BREMSSTRAHLUNG

\* S. JADACH ET AL., P.L. B377, 168 (1996)



⇒  $O(\alpha)$  CORR. TO  $\bar{\beta}_1$ , EXACT

\* FOR COMPARISON, WE ALSO IMPLEMENT  
TO RESULT OF ARBUZOV ET AL. (NPB485,457(1997))  
WHICH IS SUPPOSED TO INCLUDE  
 $O(\alpha^2 L)$ , NLLB

\* WE ALSO IMPLEMENT A SIMPLIFIED  
FORM OF OUR EXACT RESULT BASED  
ON A SOFT  $\gamma$  ANSATZ, ANSATZ(SOFT)

⇒ RESULTS IN FIGS. 1 AND 2

# LEPI




Figure 1

## LEP 1




Figure 2

$$\Rightarrow \text{FOR } 0.2 \lesssim 1 - z_{\text{min}} \lesssim 1, \quad \left. \frac{\Delta\sigma}{\sigma} \right|_{O(\alpha^2 L \bar{p}_1)} \lesssim 0.02\%$$

- EXACT RESULTS ON  $2\gamma$  BREMSSTRAHLUNG:  
TECHNICAL PRECISION CHECK
- \* WE USE EXACT RESULTS ON  $2\gamma$   
BREMSS. FOR  $\bar{\beta}_2$  (S. JADACH ET AL.,  
P.R.D41, 2682(1993); ibid. D42, 2977(1990))
- \* WE REALIZE  $2\gamma$  PHASE SPACE  
VIA AN INDEPENDENT 'TEST' MC  
FOR TECHNICAL PRECISION CHECK  
OF  $\bar{\beta}_2$
- ⇒ BY PRODUCT: CROSS CHECK PHYSICAL  
PRECISION OF  $\bar{\beta}_2$  AS WELL
- ⇒ RESULTS IN FIG. 3

## LEP1




Figure 3

$$\Rightarrow \frac{\Delta\sigma_2}{\sigma_2} \Big|_{(\mathcal{O}(\alpha^2) \bar{\beta}_2; \text{TECH. PREC.})} \leq 0.003\%$$

$$\frac{\Delta\sigma_2}{\sigma_2} \Big|_{(\mathcal{O}(\alpha^2) \bar{\beta}_2; \text{PHYS. PREC.})} \leq 0.012\%$$

(9)

• EXACT 2-LOOP VIRTUAL CORRECTION

\* WE ANALYTICALLY CONTINUE THE WORK OF BERENDS ET AL. (NP.B297, 429(1988)) TO  $t$ -CHANNEL

$$\Rightarrow \Delta \bar{\beta}_0 / \bar{\beta}_0 \Big|_{\text{Born}} = \left(\frac{\alpha}{\pi}\right)^2 L \left(6 + 6\zeta(3) - \frac{45}{8} - \frac{\pi^2}{2}\right) + \left(\frac{\alpha}{\pi}\right)^2 \left(6 - 9\zeta(3) + \left(\frac{17}{8} - 2\ln 2\right)\pi^2 - \frac{8}{45}\pi^4\right)$$

(1)

$$\Rightarrow \frac{\Delta \sigma_2}{\sigma_2} \Big|_{(\mathcal{O}(\alpha^2) \bar{\beta}_0)} \leq .014\%$$

# NEW ERROR ANALYSIS FOR BHLUMI 4.04

\* USING THE RESULTS JUST DERIVED  
WE GET

$$\frac{\Delta \sigma_L}{\sigma_L} \Big|_{O(\alpha^2)\text{-PHOTONIC}} = .027\%$$

⇒

| < JULY, 1998 > |

Type of correction/error	LEP1		LEP2
	Past	Present	Present
(a) Missing photonic $O(\alpha^2 L)$	0.15%	0.027%	0.04%
(b) Missing photonic $O(\alpha^3 L^3)$	0.008%	0.015%	0.03%
(c) Vacuum polarization	0.05%	0.04%	0.10%
(d) Light pairs	0.01%	0.03%	0.05%
(e) Z-exchange	0.03%	0.015%	0.0%
Total	0.16%	0.061%	0.122%

Table 1: Summary of the total (physical+technical) theoretical uncertainty for a typical calorimetric detector. For LEP1, the above estimate is valid for the angular range  $1^\circ$ - $3^\circ$ ; for LEP2 it covers energies up to 176 GeV, and angular range =  $[1^\circ, 3^\circ]$  (see the text for further comments).

NOTE: EXTENSION TO 200 GeV AND  
 $3^\circ$ - $6^\circ$  IN PROGRESS

(11)

• CONCLUSIONS

- ⊗ NEW ANALYSIS OF ERRORS FOR BHLUMI 4.04 COMPLETED

$O(\alpha^2 L)$  PHOTONIC ERROR REDUCED TO .027% AT LEP1

$$\Rightarrow \left. \frac{\Delta\sigma_L}{\sigma_L} \right|_{TH} = .061\% \text{ LEP1}$$

- ⊗ FOR LEP2, UP TO 176 GeV,

$$\left. \frac{\Delta\sigma_L}{\sigma_L} \right|_{TH} = .122\%$$

- ⊗ EXTENSION TO 200 GeV IN PROGRESS (AS WELL AS EXTENSION TO  $3^\circ - 6^\circ$ )

- ⊗ PREDICTIONS OF BHLUMI 4.04 UNCHANGED